INDEX TO VOLUME 72

New names and new combinations are indicated by **bold face** type

Absidia orchidis, 166; *ramosus*, 166
Acacia modesta, 399
Acenaphthene, effect of on plant tissue, 567–570
Acer pseudoplatanus, 151
Achlya flagellata, 48, 51
Achorion schoenleini, 171
Achras balata, 554; *calcicola*, 561; *chicle*, 561; *duplicata*, 554, 555; *sideroxylon*, 556; *zapota*, 551, 560; *zapotilla*, 551, 560
Achyranthus aspera, 219
Acrostichum alpinum 404; *maranteae*, 419; *septentrionale*, 416; *thelypteris*, 408; *vellea*, 419
Adanson, 52, 60–62
Adiantum capillus-veneris, 402, 421; *caudatum*, 403, 421; *pedatum*, 402, 421, 497, 501; *venustum*, 421, 422
Aecidium desmium, 206; **entebbense**, 221; *evansii*, 221; *phyllanthi*, 207, 221; *tubulosum*, 213; *vanderystianum*, 222; *vernoniae-podocomae*, 222
Aerobacter aerogenes, 158, 161
Agraulis brevifolia, 544
Agrostis, a new section *Microphyllae* in, 541; *alopecuroides*, 544; *ampla*, 542–544, 546, 547; f. **monolepis**, 547; *californica*, 542, 544; *exarata*, 541, 542, 544, 546; var. *ampla*, 542; var. *californica*, 542; var. *monolepis*, 541, 546; var. *pacifica*, 542; *hendersonii*, 541, 547; *inflata*, 543, 544, 546; **kennedyana**, 543, 544, 547; *microphylla*, 543, 544, 546; var. **hendersonii**, 543, 544, 547; var. **intermedia**, 543, 544, 547; *rossae*, 542
Aira hawaiiensis, 23; f. *depauperata*, 23; f. *haleakalensis*, 23; *praecox*, 23
Albizzia grandibracteata, 215; *lebbek*, 216; *procera*, 216
Alcaligenes fecalis, 158, 161, 163
Alcornea cordifolia, 205
Alectoria chalybeiformis, 163
Allium cepa, 570–573; *fistulosum*, 571
Amblystegiella confervoides, 516; *subtilis*, 516
Amblystegium juratzkanum, 516; *serpens*, 516; *varium*, 516
American botanical literature, index to, 114, 224, 335, 427, 483, 575
Amphilophis insculpta, 209
Anatomy of peach and cherry phloem, 137
Andreaea rothia, 511
ANDREWS, HENRY N., Cytoplasmic hybrids in *Penicillium notatum*, 361
Andropogon dummeri, 209
Angiopsora hansfordii, 206
Anisolpidium ectocarpii, 47
Anogramma leptophylla, 402
Anomodon attenuatus, 515; *minor*, 515; *rostratus*, 515; *rugelii*, 515; *tristis*, 515; *viticulosus*, 515
Antibacterial activity of fungi, 163
Antibiotic activity of lichens, 157
Antidesma venosa, 207
Arcytophyllum aristatum, 299; *capitatum*, 299; *thymifolium*, 299
Aristolochia reticulata, 270; *serpentaria*, 270
Armillaria mellea, 169
Artemisia maritima, 422
Arundinella nepalensis, 218
Ashbya gossypii, 166
Aspidium angulare, 412; *aristatum*, 413; *bakerianum*, 412; *brunonianum*, 407; *caryotideum*, 413; *drenatum*, 413; *filix-mas*, 405; *fimbriatum*, 415; *goggilodus*, 22; *gongylodes*, 22; *luctuosum*, 412; *marginatum*, 406; *rufobarbatum*, 412; *squarrosum*, 412; *tsus-simense*, 412; *varians*, 417
Asplenium acrostichoides, 414; *adiantum-nigrum*, 417, 420; *alternans*, 417; *ceterach*, 418; *filix-femina* var. *attenuatum*, 415; var. *dentigerum*, 415; var. *polysporum*, 415; var. *retusum*, 415; *fimbriatum*, 415; *fontanum*, 402, 417; *latifolium*, 426; *mackinnoni*, 414; *maximum*, 426; *mcdonelli*, 414; *pekinense*, 417; *polypodioides*, 415; *rupicola*, 414; *ruta-muraria*, 402, 416; *sarelii*, 417; *saulii*, 417; var. *pekinense*, 417; *schimperi*, 415; *septentrionale*, 402, 416; *squamigerum*, 416; *tenellum*, 414; *thelypteroides*, 414; *trichomanes*, 402, 416; *varians*, 417; *viride*, 402, 416
Athyrium acrostichoides, 402, 414; *crenatum*, 413, 416; var. *glabratum*, 416; *filix-femina*, 402, 414, 415; var. *dentigerum*, 411, 415; var. *retusum*, 415; *fimbriatum*, 415; *japonicum*, 415; *mackinnoni*, 414; *maximum*, 426; *mcdonelli*, 414, 416; *nigripes*, 414; *pectinatum*, 414; *rupicola*, 414; *schimperi*, 415; *setiferum*, 401, 414; *thelypteroides*, 414

Atrichum angustatum, 519; var. *plurilamellatum*, 519; *macmillani*, 519; *undulatum*, 519
Aulacomnium heterostichum, 513; *palustre*, 514

Bacillus cereus, 158; *mycoides*, 158, 161; *subtilis*, 158–164
Bacterium tumefaciens, 565, 566
Bacomyces aeruginosa, 163; *placophyllus*, 163; *roseus*, 163
BALDWIN, J. T., JR., Chromosomes of Cruciferae—II. Cytogeography of Leavenworthia, 367
Barbilophozia barbata, 508, 509
Barbula unguiculata, 512
BARRETT, MARY F., *Ficus altissima*, 394
Bartramia pomiformis, 514
Bassia albescens, 559
Bauhinia fassoglensis, 208
BEETLE, ALAN A., A new section *Microphyllae* in *Agrostis*, 541
Bellevalia romana, 573
BENTLEY, NANCY J., Glandular leaf hairs of oriental tobacco, 345
Betula utilis, 399
Blakea alternifolia, 390; *andreana*, 387; *bracteata*, 386; *brunnea*, 387; *calycosa*, 386; **calyptrata**, 388; var. **latior**, 388; **clusiifolia**, 387; **cuatrecasii**, 387; **florifera**, 391; *glandulosa*, 386; *intercepta*, 387; **paleacea**, 389, 390; **pilosa**, 391; **platypoda**, 390; *podagrica*, 388; *pyxidanthus*, 388; **stellaris**, 390; *subconnata*, 388, 390; var. **obtusa**, 388; **venosa**, 389, 390
Blechnum radicans, 418
Borreria laevis, 300; *ocymoides*, 300
Botanical literature, index to American, 114, 224, 335, 427, 483, 575
Bothriochloa decipiens, 211
Botrychium, the shoot of, interpreted as a series of dichotomies, 491; *boreale*, 491, 493; *chamaeconium*, 461; *dissectum*, 492, 494, 496, 500, 501; var. *obliquum*, 458, 492; *lanceolatum* var. *angustisegmentum*, 491, 494, 502; *lanuginosum*, 425, 492, 497, 498, 500; variations and origins of, 454–462; *lunaria*, 425, 491–494, 497; var. *minganense*, 492; *matricariaefolium*, 493, 494, 498, 500; subsp. *typicum*, 492; *multifidum*, 492, 494, 496, 500, 502; subsp. *silaifolium*, 492; subsp. *typicum*, 492, 496; *pumicola*, 492; *simplex*, 492; var. *compositum*, 492; var. *tenebrosum*, 492; *ternatum*, 502; *underwoodianum*, 492, 496; *virginianum*, 425, 454–462, 492, 496, 500, 502; subsp. *typicum*, 425; var. *lanuginosum*, 425
Brachiaria platynota, 207
Brachythecium acutum, 516; *digastrum*, 516; *oxycladon*, 516; *plumosum*, 517; *rivulare*, 517; *rutabulum*, 517; *salebrosum*, 517; *velutinum*, 508, 517
Brassica alboglabra, 569
Brotherella delicatula, 518; *recurvans*, 518
Bryhnia graminicolor, 517; *novae-angliae*, 517
Bryophytes of Shenandoah National Park, Virginia, 506
Bryum argenteum, 513; *bimum*, 513; *caespitosum*, 513; *capillare*, 513
Bumelia conferta, 558
BURKHOLDER, PAUL R., Further studies on the antibiotic activity of lichens, 157

Calyptrella gracilis, 473; **littoralis**, 473; *robusta*, 473; *tristis*, 473
CAMP, W. H., *Vaccinium* hybrids and the development of new horticultural material, 1
Campbellosphaera obversa, 131, 132, 134
Campylium chrysophyllum, 516; *hispidulum*, 516; *polygamum*, 516; *radicale*, 516
Candolle, A. P. de, 52, 68, 69
Cardamine uniflora, 367, 369
Cassia sieberiana, 216
Cephaëlis axillaris, 300; *jamesonii*, 296, 300, 466, 470; *tomentosa*, 300
Cephalosporium acremonium, 166
Cephalozia connivens, 509
Cephaloziella hampeana, 507, 509
Ceratodon purpureus, 511
Ceratostomella fimbriata, 166; *ips*, 166; *microspora*, 166; *minuta*, 166; *montium*, 166; *multiannulata*, 166; *obscura*, 166; *paradoxa*, 166; *penicillata*, 166; *piceaperda*, 166; *pilifera*, 166; *pini*, 166; *pseudotsugae*, 166; *radicicola*, 166; *rostrocylindrica*, 166, 179; *stenoceras*, 166; *ulmi*, 166, 179
Cerotelium deightonii, 207; *dicentrae*, 207; *gossypii*, 206
Ceterach dalhousiae, 417, 418, 420; *officinarum*, 402, 418, 420
Cetraria cucullata, 161; *juniperina*, 163; *islandica*, 163; *nivalis*, 161; *oakesiana*, 163
Chalaropsis thielavioides, 166
Chamberlainia acuminata, 517; *biventrosa*, 508, 517; *cyrtophylla*, 517
Cheilanthes albomarginata, 420; *dalhousiae*, 420; *farinosa*, 421; var. *albomarginata*, 420; var. *dalhousiae*, 420; *fragrans*, 420;

nitidula, 419; *persica*, 420; *pteridioides*, 402, 420; *subvillosa*, 419, 420; *szovitzii*, 420
Chiloscyphus pallescens, 508, 509; *polyanthus*, 508, 509; *rivularis*, 508, 509
Chomelia barbellata, 300
Chromosomes of Cruciferae—II. Cytogeography of *Leavenworthia*, 567
CHRYSLER, M. A., The shoot of *Botrychium* interpreted as a series of dichotomies, 491; The variations and origin of *Botrychium lanuginosum*, 454
Chrysophyllum, The North American species of, 191–204; *angustifolium*, 192, 201, 203; *argenteum*, 193; var. *sphaerocarpum*, 193; *bicolor*, 193, 197; *brachycalyx*, 200; *brachystylum*, 201; **brenesii**, 193, 196, 197; *caeruleum*, 193; *cainito*, 192, 198; var. *pomiferum*, 198; *claraense*, 192, 199; *eggersii*, 197; *glabrum*, 193; *gonavense*, 200; *heterochroum*, 201; **hirsutum**, 192, 198; *mexicanum*, 193, 202, 203; var. **politum**, 202, 203; var. **typicum**, 202; *miragoaneum*, 200; *montanum*, 203; *olivaeforme*, 192, 199–201, 203, 204; var. *pallescens*, 200; var. **picardae**, 200; var. *platyphyllum*, 200; var. **typicum**, 200; *pallescens*, 200; *panamense*, 193, 197; var. *macrophyllum*, 196, 197; *pauciflorum*, 193, 197; var. *krugii*, 195; var. *nervosum*, 195; *picardae*, 201; *platyphyllum*, 200; *tepicense*, 192, 198
Chusquea scandens, 470
Cinchona corymbosa, 468; *lanceolata*, 467; *officinalis*, 296, 301, 464; *pitayensis*, 296, 301; discovery and distribution of in Ecuador, 464–471; *pubescens*, 301, 305, 464, 468, 469
Cirriphyllum boscii, 517
Citrus nobilis, 569
Cladonia alpestris, 160; *amaurocraea*, 160; *apodocarpa*, 163; *atlantica*, 160; *basilaris*, 160; *borbonica*, 160, 163; *boryi*, 160; *brevis*, 160; *caespiticia*, 160, 162, 163; *capitata*, 160, 162, 163; *caroliniana*, 160; *chlorophaea*, 163; *clavulifera*, 163; *coniocraea*, 160, 164; *conista*, 163; *cristatella*, 160–164; *cryptochlorophaea*, 160, 164; *cyanipes*, 160; *deformis*, 160; *delicata*, 163; *fimbriata*, 160; *furcata*, 163; *glauca*, 160, 163; *gonecha*, 160; *gracilis* var. *choralia*, 160; var. *dilatata*, 163; var. *elongata*, 163; *grayi*, 160, 162–164; *incrassata*, 160; *macilenta*, 160; *mateocyatha*, 163; *meroclorophaea*, 160; *mitis*, 160, 161, 164; *nemoxyna*, 160; *ochrochlora*, 163; *papillaria*, 160; *pityrea*, 160; *pleurota*, 160, 161; *pyxidata*, 160; *rangiferina*, 163, 164; *robbinsii*, 160; *santensis*, 163; *scabriuscula*, 163; *squamosa*, 160; var. *denticollis*, 163; *strepsilis*, 160, 162, 163; *subcaryosa*, 163; *submitis*, 160; *subtenuis*, 160; *sylvatica*, 160, 161; *turgida*, 163; *uncialis*, 160, 161; *verticillata*, 163
Claudopus nidulans, 172, 179
Claviceps purpurea, 163
Clidemia crenulata, 479; *epiphytica*, 479; *novemnervia*, 477; **panamensis**, 478; *petiolata*, 479; *ramiflora* var. **colombiana**, 478; **vallicola**, 477
Climacium americanum, 514; *dendroides*, 514; *kindbergii*, 514
Clitocybe illudens, 169, 170; *investiens*, 169; *tabescens*, 169
Clostridium welchii, 158, 163
Cnidoscolus, the genus in Mexico, 319; *aconitifolius*, 319–322, 331; **albidus**, 320; *angustidens*, 33, 326, 327; **armatus**, 320; *calyculatus*, 326; **chaya**, 321, 322; *chayamansa*, 32, 322; *elasticus*, 319, 320, 322, 323, 328; *hernandezii*, 323, 324, 333; **jurgensenii**, 324; **liebmannii**, 324; **macrandrus**, 324, 325; *multilobus*, 324, 325, 327, 329, 334; **orbiculatus**, 325, 326; **parviflorus**, 325, 326; *pringlei*, 326, 327; **rostratus**, 328; *souzae*, 328; **spinosus**, 320, 329, 330; **tenuilobus**, 319, 330; *tepiquensis*, 320, 323, 330–332; **tomentosus**, 332, 333; *tubulosus*, 32, 41, 320; **velutinus**, 325, 333, 334
Coccocypselum condalia, 302; *umbellatum*, 302
Coffea arabica, 302
Colchicine, effect of on plant tissue, 563
Collybia radicata, 168; *velutipes*, 168
Commelina vogelii, 208
Condaminea corymbosa, 302
Coniogramme affinis, 419; *fraxinea*, 419
Coniophora cerebella, 172; *suffocata*, 172, 179
Conocephalum conicum, 511
Conostegia cuatrecasii, 473; *poeppigii*, 474; **rubiginosa**, 473
Coprinus radians, 168
Cornus alba var. *flaviramea*, 223; *asperifolia*, 223; *disciflora*, 223; var. *floccosa*, 223; f. **floccosa**, 223; *drummondi*, 223; f. **priceae**, 223; *florida*, 223; subsp. **urbiniana**, 223; var. *urbiniana*, 223; *priceae*, 223; *stolonifera*, 223; var. *flaviramea*, 223; f. *baileyi*, 223; f. **flaviramea**, 223; *urbiniana*, 223

Corticium abeuns, 170, 172, 180, 182–185, 187–189; *caeruleum*, 169; *confluens*, 169; *conigenum*, 172; *cremoricolor*, 168; *galactinum*, 169; *hydnans*, 172, 180, 185, 186; *lividum*, 170, 172, 179; *roseum*, 169, 172; *salmonicolor*, 168
Corynula pilosa, 302
Cotton, effect of on the germination of *Phycomyces* spores, 76
CROIZAT, LEON, "*Euphorbia chamaesyce*" in the United States, 312; History and nomenclature of the higher units of classification, 52
CRONQUIST, ARTHUR, Studies in the Sapotaceae—I. The North American species of *Chrysophyllum*, 191; IV. The North American species of *Manilkara*, 550
Crossopsora antidesmiae-dioicae, 207
Cryptogramma brunnoniana, 418; *crispa*, 418; *stelleri*, 402, 418
CUMMINS, GEORGE B., Descriptions of tropical rusts—VII, 205
Cyanococcus cuthbertii, 5, 12
Cyclophorus porosus, 401, 424
Cyclosorus gongylodes, 22
Cycnoches chlorochilon, 214
Cymbopogon confertiflorus, 209; *excavatus*, 209; *giganteus*, 209; *martini*, 209
Cyrtomium caryotideum, 403, 413
Cystopteris fragilis, 399, 402, 404; *montana*, 402, 404; *retusa*, 415
Cytidia salicina, 168
Cytoplasmic hybrids in *Penicillium notatum*, 361

Dactylina arctica, 163
Daedalia ambigua, 168; *confragosa*, 168; *heteromorpha*, 172; *juniperina*, 172; *quercina*, 172; *unicolor*, 168
DARROW, GEORGE M., *Vaccinium* hybrids and the development of new horticultural material, 1
Datura stramonium, 254
Davallia pulchra, 413
DAVIDSON, ROSS W., A survey of some wood-destroying and other fungi for antibacterial activity, 165
Dermatocarpon miniatum, 163
Dermatophytes, 171, 178
Deschampsia australis, 23; **hawaiiensis**, 23; f. **depauperata**, 23; f. **haleakalensis**, 23
Descriptions of tropical rusts—VII, 205
Dicranella heteromalla, 511; *varia*, 508, 511
Dicrano-weisia crispula, 508, 512
Dicranum bonjeani, 512; *flagellare*, 512; *fulvum*, 512; *fuscescens*, 512; *montanum*, 512; *rugosum*, 512; *scoparium*, 512
Dioclea reflexa, 219
Diplazium fraxineum, 419; *japonicum*, 415; *maximum*, 401, 426; *polypodioides*, 415, 426; *squamigerum*, 413, 416
Diphysium foliosum, 519
Diplococcus pneumoniae, 158, 161
Diplophyllum apiculatum, 510
Discovery and distribution of *Cinchona pitayensis* in Ecuador, 464
Ditrichum lineare, 508, 511; *pallidum*, 511; *pusillum*, 511
Drosophila melanogaster, 235
Drummondia prorepens, 514
Dryopteris, *Deschampsia*, *Portulaca*, *Lupinus*, *Fagara*, *Stenogyne*, and *Dubautia*. Hawaiian plant studies 12, 22
Dryopteris africana, 401, 409; *barbigera*, 407, 411; var. *falconeri*, 407; *blanfordii*, 405, 406; *brunnea*, 409, 426; *brunoniana*, 407, 411; *cochleata*, 406; var. *schimperiana*, 406; *crenata*, 407; *dentata*, 403, 409; *erubescens*, 408; *filix-mas*, 402, 405, 406; var. *odontoloma*, 405; *goggylodus*, 22, 23; *gongylodes*, 22, 23; *hexagonoptera*, 401, 411; *hirtipes*, 401; var. *atrata*, 408; *laterepens*, 400, 409, 426; *levingei*, 409; *linnaeana*, 402, 410, 411; *ludens*, 403; *macrocarpa*, 401, 406; *marginata*, 401, 405, 406; *moulmeinensis*, 408; *odontoloma*, 405; *paleacea*, 405; *parasitica*, 409; *penangiana*, 408; *phegopteris*, 402, 410; *prolifera*, 401, 403, 409; *punctata*, 411; *ramosa*, 405, 406; *rampans*, 403, 408; *repens*, 401, 408; *robertiana*, 402, 410, 411; *rosthornii*, 404, 405; *schimperiana*, 401, 406; *serratodentata*, 406; *subpubescens*, 409; *thelypteris*, 402, 408
Dubautia, 22; *plantaginea*, 26–28; var. *angustifolia*, 26, 28, 29; var. *chamissonii*, 26, 30; var. **glandulosa**, 30; var. **paucaensis**, 30; var. *strigosa*, 30; var. **typica**, 26, 30
DWYER, JOHN D., The taxonomy of the genus *Sauvagesia* (Ochnaceae), 521

Echinodonta tinctorum, 169
Ecuador, *Cinchona pitayensis* in, 464; Rubiaceae from, 295
Effect of cotton on the germination of *Phycomyces* spores, 76
Effect of colchicine and acenaphthene in combination with x-rays on plant tissue —I. Introduction, 563

Elaeagnus utilis, 302
Encalypta streptocarpa, 512
Endoconidiophora adiposa, 166; *caerulescens*, 166
Entodon brevisetis, 517; *cladorrhizans*, 517; *seductrix*, 517
Epidermophyton inguinale, 171, 178
Equisetum arvense, 425; *debile*, 425, 426; *ramosissimum*, 425, 426
Escherichia coli, 158, 161, 163, 165–189
Espeletia hartwigiana, 466
Estimates of the minimum numbers of genes differentiating several taxonomic categories in the genus *Solidago*, 235
"Euphorbia chamaesyce" in the United States, 312
Euphorbia callitrichoides, 317; *chamaesyce*, 312–317; var. *canescens*, 315; *humifusa*, 316; *milii*, 313; *portulacoides*, 317; *prostata*, 312–317; *thymifolia*, 315
Eurhynchium hians, 517; *rusciforme*, 517; *serrulatum*, 517; *strigosum*, 517; var. *praecox*, 517
EVANS, ALEXANDER W., Further studies on the antibiotic activity of lichens, 157
Evernia vulpina, 163
Exidia glandulosa, 168

Fagara, 22; *dipetala* var. *geminicarpa*, 24; *mauiensis* var. **anceps f. petiolulata**, 25
Faramea coerulescens, 302; *maynensis*, 302; *oblongifolia*, 302
FASSETT, NORMAN C., *Juniperus virginiana*, *J. horizontalis*, and *J. scopulorum*—III. Possible hybridization of *J. horizontalis* and *J. scopulorum*, 42; IV. Hybrid swarms of *J. virginiana* and *J. horizontalis*, 379; V. Taxonomic treatment, 480
Favolus canadensis, 168
Ferns of Kashmir, 399
Ficus altissima, 394–398; var. *fergusonii*, 397, 398; var. *laccifera*, 394, 397; f. *laccifera*, 394; *bengalensis*, 394, 395, 397, 398; *capensis*, 207; *indica*, 394, 397; *laccifera*, 394, 396–398; *peguensis*, 394, 397, 398; *varenga*, 394, 397, 398; *variegatus*, 207
Filix montana, 404
Fissidens adiantoides, 511; *cristatus*, 511; *minutulus*, 511; *osmundioides*, 511; *subbasilaris*, 511; *taxifolius*, 511; *viridulus*, 508
Fistulina hepatica, 172
Flammula alnicola, 172, 179
Fomes annosus, 172, 180; *aplanatus*, 171, 172; *calkinsii*, 168; *clelandii*, 169; *conchatus*, 169; *connatus*, 168; *densus*, 168; *everhartii*, 168; *extensus*, 169; *ferruginosus*, 172; *fometarius*, 168; *fraxineus*, 168; *fraxinophilus*, 172, 173; *fulvus*, 169, 170; *geotropus*, 169; *igniarius*, 168; var. *laevigatus*, 168; var. *populinus*, 169; *juniperinus*, 173, 180, 187; *lobatus*, 169; *marmoratus*, 169; *meliae*, 173; *officinalis*, 173; *pini*, 173; var. *abietis*, 173; *pinicola*, 173; *pomaceus*, 169, 170; *ribis*, 173, 180–182, 187; *rimosus*, 168, 169; *robustus*, 168; var. *tsugina*, 168; *roseus*, 173; *scutellatus*, 169; *subroseus*, 173; *tenuis*, 173; *torulosus*, 168; *ulmarius*, 179; *ungulatus*, 173
Fontinalis novae-angliae, 514
Fraxinus excelsior, 151
Frullania asagrayana, 510; *brittoniae*, 510; *eboracensis*, 510; *riparia*, 510
FULLING, EDMUND H., Thomas Jefferson. His interest in plant life as revealed in his writings—II, 248
Further studies on the antibiotic activity of lichens, 157
Fusarium avenaceum, 166

Galium aparine, 303; *canescens*, 303; *piliferum*, 303; *tetraplazium*, 303
Ganoderma sp., 173
Gastridium lendigerum, 547
Gaylussacia brachycera, 10
Genus *Cnidoscolus* in Mexico: new species and critical notes, 319
Genus *Jatropha* in America: principal intrageneric groups, 271
Gibberella saubinetti, 166; *zeae*, 166
Glandular leaf-hairs of oriental tobacco, 345
GLEASON, H. A., On *Blakea* and *Topobea*, 385; Some Melastomaceae of Colombia, 472
Gomozia granadensis, 303
Gomphosphaeria lacustris, 246
Goniophlebium lachnopus, 423; *microrhizoma*, 422; *argutum*, 423
Goniopteris lineata, 408; *prolifera*, 409
Gonzalagunia cornifolia, 303; *dependens*, 304; *runis*, 304; *soraria*, 304
GOODWIN, RICHARD H., Estimates of the minimum numbers of genes differentiating several taxonomic categories in the genus *Solidago*, 235
Gossypium acuminatum, 206
Grammitus vestita, 418
Grimmia alpicola var. *rivularis*, 512; *apocarpa*, 513; var. *conferta*, 513; var. *gracilis*, 513; *pilifera*, 513
Grosmannia serpens, 166

Guettarda hirsuta, 304; *sabiceoides*, 304
Gymnocarpium dryopteris, 410; *remotum*, 410; *robertianum*, 410, 411
Gymnogramma aurita var. *levingei*, 409; *javanica*, 418; *leptophylla*, 402; *levingei*, 409; *totta*, 409; *vestita*, 418
Gymnopteris vestita, 418

Hamelia patens, 304; *pedicellata*, 304
Hawaiian plant studies, 22
Hedwigia ciliata, 514
Hemarthria uncinata, 211
Hemidictyum ceterach, 418
Hemidiodia ocimifolia, 304
Hemionitis prolifera, 409
HERVEY, ANNETTE, A survey of some wood-destroying and other fungi for antibacterial activity, 165
Heterophyllum haldanianum, 518
Hillia parasitica, 304
Hirneola sp, 173
History and nomenclature of the higher units of classification, 52
Hoffmannia ecuatoriana, 305; *sprucei*, 305
Homomallium adnatum, 518
Hyacinthus orientalis, 571
Hydnum abietis, 173, 180, 186; *caput-ursi*, 174; *chrysocomum*, 168; *coralloides*, 169, 170; *erinaceus*, 169, 170; *ochraceum*, 168; *pulcherrimum*, 168; *scheidermayeri*, 174, 179; *septentrionale*, 174, 180, 184, 186
Hydrastis canadensis, 270
Hygroamblystegium fluviatile, 516; var. *ovatum*, 516; *irriguum*, 516; var. *spinifolium*, 516; *orthocladon*, 516
Hygrohypnum dilatatum, 516; *eugyrium*, 516
Hylocomnium brevirostre, 519; *splendens*, 519; *umbratum*, 508
Hymenochaete agglutinans, 169; *corrugata*, 174, 179; *curtisii*, 169; *rubiginosa*, 169, 174, 179; *tabacina*, 174
Hyparrhenia rufa, 209
Hypnum arcuatum, 518; *cupressiforme*, 518; *curvifolium*, 518; *fertile*, 518; *imponens*, 518; *molluscum*, 518; *pallescens*, 508, 518; *reptile*, 518
Hypholoma sublateritium, 174
Hypolepis punctata, 411

Index to American botanical literature, 114, 224, 335, 427, 483, 575
Indigofera arrecta, 216
Inheritance of orange and yellow flower color in *Nemesia strumosa*, 435
Inula acervata, 221
Irpex cinnamomeus, 174; *mollis*, 174, 180, 186, 187, 189
Isertia alba, 305; *pittieri*, 305
Ithyphallus sp., 174

Jamesoniella autumnalis, 510
Jasmineum pauciflorum, 218
Jatropha, The genus in America, 271; *acerifolia*, 292; *aconitifolia*, 319; *acuminata*, 274, 292; *alamani*, 33, 35, 288, 289, 292; *andrieuxii*, 276, 278, 283, 284, 292; *angustifolia*, 274, 292, 293; *antisiphilitica*, 292; *arizonica*, 292; *augusti*, 276–278, 292; *berlandieri*, 32, 276, 277, 292; *berterii*, 292; *bornmülleri*, 280, 292; *brachypoda*, 280, 281, 292; *breviloba*, 280, 292; *canescens*, 289, 292; *capensis*, 274; *cardiophylla*, 33, 288, 291, 292; *cathartica*, 32, 272, 277, 292; *catingae*, 291, 292; *cercidiphylla*, 292; *ciliata*, 35, 36, 41, 276–279, 288, 289, 292, 293; *cinerea*, 33, 36, 286–289, 292, 293; *clavuligera*, 280, 282, 293; *coccinea*, 293; *cordata*, 33, 35, 37, 272, 288, 290, 291, 293; *cuneata*, 36, 286, 287, 293; *cuneifolia*, 293; *curcas*, 33, 35, 271–273, 278, 283–285, 292–294; var. *rufa*, 272, 284; *decipiens*, 293; *dioica*, 36, 37, 41, 286, 290–293; var. **graminea**, 36, 39, 287; var. **sessiliflora**, 36, 37, 41, 287; *dissecta*, 280, 281, 293; *divaricata*, 273, 286, 293; *divergens*, 293; *diversifolia*, 274, 275, 293; *edulis*, 36, 37, 41, 292, 293; *elegans*, 293; *elliptica*, 280, 281, 293; *excisa*, 272, 280, 293; *flabellifolia*, 293; *flavovirens*, 280, 281, 293; *fremontioides*, 285, 286, 293; *gaumeri*, 275, 285, 286, 293; *glauca*, 293; *glaucovirens*, 274, 275, 293; *gossypifolia*, 40, 273, 275, 278, 279, 281, 282, 293, 294; var. *elegans*, 32; *grandifrons*, 35, 41, 293; *grossidentata*, 276, 277, 293; *guaraniticia*, 280, 293; *harmsiana*, 290, 293; *hastata*, 33, 273–275, 293; *hernandiaefolia*, 33, 286, 293, 294; *hernandifolia*, 273; *heterophylla*, 33, 41, 293; *hieronymi*, 276, 277, 293; *hippocastanifolia*, 281, 293; *hoffmanniae*, 35, 276, 277, 293; *humboldtiana*, 35, 41, 272, 277, 293, 294; *induta*, 280, 281, 293; *integerrima*, 33, 272, 274, 275, 278, 292–294; *intercedens*, 281, 293; *intermedia*, 280, 293; *isabelli*, 280, 281, 292, 293; var. *antisiphilitica*, 280; *jacquini*, 293; *jurgensenii*, 319, 324; *katharinae*, 280, 293; *lacerti*, 293; *liebmannii*, 319, 324; *loeflingii*, 293; *longipedunculata*, 35, 276, 293; *luxurians*, 293; *macrantha*, 271,

275–277, 279, 293; *macrocarpa*, 276, 277, 279, 293; *macrorhiza*, 279, 281, 292, 293; var. *septemfida*, 292; *malacophylla*, 287, 288, 290; *malmeana*, 281, 293; *martinsii*, 276, 293; *minor*, 275, 293; *mitis*, 293; *mollissima*, 279, 293, 294; *moluensis*, 33, 41, 275, 293; *multifida*, 32, 273, 276, 277, 293; *multiflora*, 276, 293; *multiloba*, 324; *mutabilis*, 276, 277, 293; *neopauciflora*, 37, 286, 287, 290, 293, 294; *nudicaulis*, 276, 277, 293; *octandra*, 41; *officinalis*, 293; *olivacea*, 32, 35, 41, 293; *opifera*, 293; *ortygae*, 286, 288, 290, 293; *pachypoda*, 276, 277, 279, 293; *palmata*, 39, 41; *palmatifolia*, 291, 293; *palustris*, 293; *panduraefolia*, 274, 275, 294; *papyrifera*, 276, 277, 294; *pauciflora*, 274, 275, 294; *paxii*, 274, 293, 294; *pedantipartita*, 281, 282, 294; *peiranoi*, 280, 294; *peltata*, 34, 41, 276, 277, 294; *platanifolia*, 278, 290, 294; *platyphylla*, 32, 34, 286, 294; *podagrica*, 32, 34, 276, 277, 294; *pohliana*, 276, 277, 279, 294; *portoricensis*, 294; *prunifolia*, 274; *pseudocurcas*, 282, 284, 294; *puncticulata*, 280, 294; *purpurea*, 280, 281, 294; *quinqueloba*, 40, 41; *ribifolia*, 280, 281, 294; *ricinifolia*, 276, 277, 279, 294; *rigidifolia*, 280, 294; *riojae*, 286, 288, 291, 294; *robertii*, 277, 294; *rufescens*, 287, 294; *spathulata*, 36, 37, 41, 290, 294; var. *genuina*, 37; var. *sessiliflora*, 33, 37; *standleyi*, 286, 288, 291, 294; *staphysagrifolia*, 294; *stigmatosa*, 291, 294; *sympetala*, 286, 287, 291, 294; *tacumbensis*, 281, 294; *tepiquensis*, 331; *thyrsantha*, 280, 281, 294; *transiens*, 280, 281, 294; *triloba*, 39–41; *tubulosa*, 40; var. *quinqueloba*, 40; var. *septemloba*, 40; *tupifolia*, 274, 275, 294; *unicostata*, 274; *variegata*, 274; *variifolia*, 274; *vernicosa*, 288, 291, 294; *villosa*, 294; *weberbaueri*, 35, 276, 277, 294; *wedeliana*, 276, 277, 279, 294; *yucatanensis*, 35, 41, 294

Jatrophas of Cervantes and of the Sessé and Mociño herbarium, 31

Jaume-St. Hilaire, 52, 69–71

Jefferson, Thomas, 248

Jeffersonia, 269; *binata*, 270; *diphylla*, 270

Joosia pulcherrima, 305

Jubula pennsylvanica, 510

Juglans alba, 252; *pecan*, 253; *paccan*, 253

Juniperus communis var. *depressa*, 43; *horizontalis*, 42, 43, 45, 46, 379–383, 480–482; *lucayana*, 482; *sabina* var. *procumbens*, 46; *scopulorum*, 42–46, 379–384, 480–482; var. **columnaris**, 480–482; var. **patens**, 44–46, 480–482; *silicicola*, 482; *virginiana*, 45, 46, 379–384, 480–482; var. **ambigens**, 380–383, 480–482; var. *crebra*, 480–482

Juniperus virginiana, *J. horizontalis*, and *J. scopulorum*—III. Possible hybridization of *J. horizontalis* and *J. scopulorum*, 42; IV. Hybrid swarms of *J. virginiana* and *J. horizontalis*, 379; V. Taxonomic treatment, 480

Jussieu, Antoine Laurent de, 64, 65; Bernard de, 65

KARLING, JOHN S., *Rhizidiomyces hirsutus* sp. nov., a hairy anisochytrid from Brazil, 47

Kashmir, The ferns of, 399

Kuehneola desmium, 206; *gossypii*, 206

Lactobacillus casei, 158, 163

Ladenbergia macrocarpa, 305; *magnifolia*, 305

Lamarck, 52, 63, 64

Lastrea atrata, 408; *barbigera*, 407; *brunoniana*, 407; *crenata*, 407; *falconeri*, 407; *filix-mas* var. *elongata*, 406; var. *odontoloma*, 406; var. *parallelogramma*, 404; var. *schimperiana*, 406; var. *serrato-dentata*, 406; *hirtipes*, 408; *odontoloma*, 405; *thelypteris*, 408

Leavenworthia, cytogeography of, 367; *aurea*, 367–369, 373–377; *michauxii*, 367, 369; *stylosa*, 368–370, 372–377; *torulosa*, 367–373, 375, 377; *uniflora*, 367, 368, 370, 373, 375–377

Leitgebia imthurnia, 528

Lejeunea cavifolia, 511; *patens*, 511

Lentinus lepidius, 169, 170; *tigrinus*, 168; *villosa*, 168

Lenzites betulina, 168; *trabea*, 168–170; *saepiaria*, 170, 174

Lepiota naucina, 174, 180, 186, 187

Lepisorus loriformis, 423

Leptodictyum trichopodium var. *kochii*, 516

Leptogramme totta, 409

Leptostegia lucida, 421

Leskea obscura, 515; *polycarpa*, 515

Leucobryum albidum, 512; *glaucum*, 512

Leucodon brachypus, 514; *julaceus* f. *flagelliferus*, 514; *sciuroides*, 514; *trichomitrium*, 515

Leucolejeunea clypeata, 511

Leucostegia pulchra, 401, 413

LEVINE, MICHAEL, The effect of colchicine and acenaphthene in combination with

x-rays on plant tissue—I. Introduction, 563
Lichens, further studies on antibiotic activity of, 157
LINDEGREN, CARL C., Cytoplasmic hybrids in *Penicillium notatum*, 361
Lindley, 52, 70
Linnaeus, 52, 54–63
Lippia asperifolia, 221; *berlandieri*, 222
Lobaria pulmonaria, 163
Lophocolea bidentata, 509; *heterophylla*, 509; *minor*, 507, 509
Loureira cuneifolia, 37; *glandulosa*, 290
LUNDELL, CYRUS LONGWORTH, The genus *Cnidoscolus* in Mexico; new species and critical notes, 319
Lupinus, 22; *albus*, 571; var. *termis*, 24
Lycoperdon gemmatum, 169; *pyriforme*, 169
Lycopersicum esculentum, 571; *pimpinellifolium*, 571
Lycopodium sanguinolentum, 426; *selago*, 426; f. *angustinum*, 426
Lygodium japonicum, 424

MA, ROBERTA, A survey of some wood-destroying and other fungi for antibacterial activity, 165
Macrocnemum sprucei, 306
Manetta echitidea, 306; *evenia*, 306; *flexilis*, 306; *pichinchensis*, 306; *racemosa*, 306; *recurva*, 306; *skutchii*, 306; *trianae*, 306
Manihot aesculifolia, 39; *fetida*, 32, 34, 40; *palmata*, 39
Manilkara, The North American species of, 550; **albescens**, 553, 558; *bidentata*, 551, 552, 554; *breviloba*, 560; *calcicola*, 561; var. *columbiana*, 561; *calderonii*, 560; *chicle*, 553, 561; *conzattii*, 560; *darienensis*, 553; *duplicata*, 554; *emarginata*, 551, 553, 556, 558; subsp. **haitensis**, 558; subsp. **jaimiqui**, 557–559; subsp. **typica**, 557; subsp. **wrightiana**, 557; **excisa**, 552, 555; *gaumeri*, 560; **gonavensis**, 553, 562; *grisebachii*, 557, 558; *howardii*, 557; *jaimiqui*, 557; **mayarensis**, 553, 558, 559; *meridionalis*, 551, 553, 560; var. *caribbensis*, 560; var. *meridionalis*, 560; *nitida*, 553; *pleeana*, 552, 554; *riedleana*, 553; *rojasii*, 560; *sideroxylon*, 553, 555, 556; *sapodilla*, 552; *spectabilis*, 552, 555; *staminodella*, 551, 553, 560; *striata*, 560; *tabogaensis*, 560; *wrightiana*, 557; *zapotilla*, 551, 553, 556, 560
Manisurus altissima, 211
Marsilia crosa, 425; *minuta*, 425; *natans*, 425
Marsupella emarginata, 510
MCVAUGH, ROGERS, The genus *Jatropha* in America: principal intrageneric groups, 271; The Jatrophas of Cervantes and of the Sessé and Mociño herbarium, 31
Melandryum viscosum, 349
Melastomaceae of Colombia, 472
Melinis tenuissima, 206
Mentha piperita, 350
Merrillosphaera carteri, 132
Merulius confluens, 169; *niveus*, 168; *tremellosus*, 168
Metzgeria conjugata, 511; *crassipilis*, 511
METZNER, JEROME, A morphological and cytological study of a new form of *Volvox*—I, 86; II, 121
Miconia axinaeoides, 476; *barbinervis*, 477; **blakeaefolia**, 475; **calcarata**, 474; **centronioides**, 475, 476; *clavistyla*, 477; *compacta*, 477; *dolichorrhyncha*, 477; *gracilis*, 476; *lanata*, 477; *marginata*, 476; *megalantha*, 476; *pilgeriana*, 477; **platypoda**, 476; *ruficalyx*, 477; *semota*, 477; *stelligera*, 477; **stipitata**, 476; **transversa**, 475
Microlepidozia sylvatica, 509
Microlepis wilfordii, 413
Microsporum canis, 178; *felineum*, 171; *gypseum*, 171
Mimena holwayi, 216
Morchella esculenta, 166, 168
Morphological and cytological study of a new form of *Volvox*—I, 86; II, 121
Mozinna canescens, 290; *cordata*, 290; *sessiliflora*, 37; *spathulata*, 36, 37, 282, 285; var. *sessiliflora*, 37
Mucuna urens, 219
Mimusops albescens, 559; *azulensis*, 559; *balata*, 554; var. *domingensis*, 553; var. *sieberi*, 553; *bidentata*, 553, 554; *camagueyensis*, 557; *darienensis*, 553; *duplicata*, 554; *excisa*, 555; *floridana*, 557; *gonavensis*, 561; *grisebachii*, 557; *jaimiqui*, 557; *jamaicensis*, 556; *kaukii*, 550; *mayarensis*, 558; *nitida*, 553; *parviflora*, 557; *pleeana*, 554; *riedleana*, 553; *sideroxylon*, 556; *sieberi*, 553, 557; *wrighteana*, 557
Mniobryum albicans, 513
Mnium affine, 513; *cuspidatum*, 513; *hornum*, 513; *marginatum*, 513; *orthorhynchum*, 508, 513; *punctatum*, 513; *stellare*, 513
Murica albescens, 559; *eyerdamii*, 559

Necker, 52, 66
Neckera pennata, 515
Nemesia compacta, 436; *grandiflora*, 436; *strumosa*, 435–452; inheritance of orange and yellow flower color in, 435; *strumosa suttoni*, 436; *versicolor*, 436
Nephrodium aristatum, 413; var. *affine*, 413; *barbigerum*, 407; *blanfordii*, 405; *brunonianum*, 407; *costatum*, 408; *falconeri*, 407; *filix-mas*, 405; var. *elongatum*, 406; var. *fibrillosum*, 404; var. *marginatum*, 406; var. *normale*, 405; var. *schimperianum*, 406; *hirtipes*, 408; *kingii*, 406; *marginatum*, 406; *molle*, 409; *odontoloma*, 405; *prolixum*, 408; *rampans*, 405; *remotum*, 405; *repens*, 408; *rigidum*, 405; *rosthornii*, 404; *schimperianum*, 406; *serrato-dentatum*, 406
Nephromopsis ciliaris, 163
Nertera depressa, 303
Neurospora crassa, 166; *sitophila*, 166
New combinations in *Cornus*, 223
New section *Microphyllae* in *Agrostis*, 541
Nicotiana, 345–360; alata, 438; glauca, 566; langsdorfii, 438
Niphobolus fissus, 424; *porosus*, 424
Notholaena marantae, 401, 402, 419; *persica*, 420; *vellea*, 401, 402, 419
Nowellia curvifolia, 509

Odontoschisma denudatum, 509
Olea cuspidata, 399
Omphalia campanella, 174
On *Blakea* and *Topobea*, 385
Onychium contiguum, 421; *japonicum* var. *lucidum*, 421; var. *multisectum*, 421; *lucidum*, 421
Ophiobolus oryzinus, 166
Ophioglossum engelmanni, 497, 501; *fibrosum*, 498; *japonicum*, 424; *vulgatum*, 425
Ophiostoma catonianum, 166
Orthotrichum ohioense, 514; *speciosum*, 514
Osmunda claytoniana, 402, 411, 424; *lunaria*, 425; *regalis*, 402; *virginiana*, 425
Ossaea resinosa, 478
Oxalis violacea, 371
Oxycoccus macrocarpus, 1

Palicourea angustifolia, 306; *balnearia*, 307; *calantha*, 307; *caldasiana*, 307; *calothyrsus*, 307; *chimboracensis*, 307; *crocea*, 307; *egena*, 307; *flavescens*, 307; *guianensis*, 307; *holmgrenii*, 307; *hospitalis*, 307; *huigrensis*, 308; *kalbreyeri*, 308; *lasiantha*, 308; *levis*, 308; *lugubris*, 308; *oreadium*, 308; *pasti*, 308
Panicum deustum, 207
Panus laevis, 168; *rudis*, 168; *stipticus*, 174; *strigosus*, 168
Paraleucobryum longifolium, 512
Parmelia caperata, 161; *cladonia*, 163; *conspersa*, 163; *physodes*, 161–164; *rudecta*, 163
Paspalum commersonii, 212; *dilatum*, 212; *orbiculare*, 212
Paxillus involutus, 174
Pelargonium zonale, 566
Pellaea gracilis, 418; *hastata*, 419; *nitidula*, 419; *stelleri*, 418
Peltigera variolosa, 163
Penicillium luteum, 361; *notatum*, 165, 179, 182–184, 188, 189, 361–365
Peniophora allescheri, 168; *cinerea*, 169, 170; *coccineofulva*, 169; *gigantea*, 168; *incarnata*, 169, 170; *nuda*, 169; *pubera*, 168
Pennisetum polystachyum, 206
Phacopsora apoda, 206; *desmium*, 206; **greweae**, 206, 210, 211; *hansfordii*, 205; *stereospermi*, 205
Phegopteris distans, 426; *erubescens*, 408; *hexagonoptera*, 411; *punctata*, 411; *robertiana*, 410; *vulgaris*, 410
Philonotis americana, 514; *fontana*, 514
Phymatodes stracheyi, 423
Phlebia strigosozonata, 180, 186
Pholiota adiposa, 166, 168; *aeruginosa*, 175; *aurivella*, 168; *limonella*, 168; *lutea*, 169; *spectabilis*, 168, 169
Phragmopyxis leonensis, 216, 217
Phycomyces blakesleeanus, 76–85, 166
Phyllanthus discoideus, 207, 221
Physcia sp., 163
Physcomitrium turbinatum, 513
Phytophthora cinnamomi, 166
Pinus excelsa, 399; *longifolia*, 399, 422
Pistia stratiotes, 87
Pisum sativum, 573
Plagiochila ludoviciana, 508, 510
Plagiothecium denticulatum, 517; *deplanatum*, 517; *elegans*, 518; *lactum*, 518; *muellerianum*, 518; *roseanum*, 518; *sylvaticum*, 518; *turfaceum*, 518
Plectocolea crenulata, 510
Pleopeltis clathrata, 424; *membranacea*, 423; *stewartii*, 423
Pleurotus corticatus, 168; *griseus*, 175, 180–189; *ostreatus*, 168; *sapidus*, 168; *ulmarius*, 168
Pleurozium schreberi, 516
Podophyllum diphyllum, 270

Pogonatum pensilvanicum, 519
Pohlia cruda, 513; *nutans*, 513
Polycodium melanocarpum, 2; *stamineum*, 2
Polypodium aculeatum, 412; *africanum*, 409; *argutum*, 401, 423; *aristatum*, 413; *brunneum*, 426; *calcareum*, 410; *clathratum*, 423, 424; *crenatum*, 407; *dentatum*, 409; *dentigerum*, 415; *distans*, 426; *erubescens*, 408; *excavatum*, 423, 424; *filix-femina*, 414; *filix-mas*, 405; *fontanum*, 417; *fragile*, 414; *fragrans*, 420; *hexagonopterum*, 411; *lachnopus*, 401, 423; *latirepens*, 400, 409; *lineare*, 402; var. *loriforme*, 423; var. *steniste*, 423; *lineatum*, 406; *lonchitis*, 411; *loriforme*, 402, 423, 424; *membranaceum*, 403, 423; *microrhizoma*, 401, 422; *montanum*, 404; *phegopteris*, 410; *porosum*, 424; *proliferum*, 409; *pteridioides*, 420; *punctatum*, 411; *robertianum*, 410; *stewartii*, 423; *stracheyi*, 401, 423
Polypogon alopecuroides, 544; *monspeliensis* var. *monolepis*, 544
Polyporus abietinus, 168; *albellus*, 175, 180; *alboluteus*, 175; *adustus*, 168; *anceps*, 168, 188; *arcularius*, 168; *balsameus*, 175; *berkeleyi*, 168; *betulinus*, 175; *biformis*, 175, 180; *borealis*, 169; *brumalis*, 168; *cinnebarinus*, 168; *circinatus*, 169; *compactus* 169; *cristatus*, 169; *croceus*, 169; *curtisii*, 169; *cuticularis*, 168; *delectans*, 175, 180; *dichrous*, 168; *distortus*, 168; *dryadeus*, 168; *dryophilus*, 168; var. *vulpinus*, 169; *durescens*, 175; *ectypus*, 168; *fibrillosus*, 168; *fissilis*, 168; *fragilis*, 175; *frondosus*, 168; *fumidiceps*, 175; *fumosus*, 168; *galactinus*, 175; *giganteus*, 175, 180, 187; *gilvus*, 168; *glomeratus*, 168; *graveolens*, 168; *guttulatus*, 175; *hirsutus*, 168; *hispidus*, 168; *hydnoides*, 168; *immitus*, 175; *lucidus*, 168; *ludovicianus*, 169; *mutabilis*, 168; *obtusus*, 175, 180; *oregonensis*, 169; *osseus*, 169; *palustris*, 176; *pargamenus*, 168, 169; *picipes*, 168; *pinsitus*, 168; *pubescens*, 168; *radiatus*, 176, 180; *resinosus*, 168, 176, 180; *robiniophilus*, 168; *rutilans*, 168; *schweinitzii*, 169, 170, 176, 180; *spraguei*, 169, 170; *spumeus*, 168; var. *malicola*, 168; *squamosus*, 168; *subcartilagineus*, 176; *subectypus*, 168; *sulphureus*, 169; *supinus*, 168; *tenuis*, 168; *texanus*, 168; *trabeus*, 176, 180; *tuberaster*, 169; *tuckahoe*, 169; *tulipiferus*, 176, 179; *varius*, 169; *velutinus*, 168; *versicolor*, 166, 168; *volvatus*, 169; *vulpinus*, 176, 179; *zonalis*, 168; *zonatus*, 168
Polystichum aculeatum, 402, 412, 413; var. *rufobarbatum*, 412; *angulare*, 412, 413; *aristatum*, 401, 413; *bakerianum*, 412; *falcatum*, 413; subsp. *caryotideum*, 413; *lachenense*, 411; *lonchitis*, 411; *prescottianum*, 411, 412; var. *bakerianum*, 412; *squarrosum*, 412; *thomsoni*, 411, 412; *tsus-simense*, 412
Polytrichum commune, 519; *juniperinum*, 519; *ohioense*, 519; *piliferum*, 508, 519
Porella pinnata, 510; *platyphylloidea*, 510
Porotrichum alleghaniense, 514
Poria ambigua, 168; *andersonii*, 169; *carbonica*, 169, 170; *cocos*, 176; *colorea*, 176; *corticola*, 176, 180, 182, 183, 187, 188; *ferrea*, 176; *ferruginea-fusca*, 176, 180, 187; *ferruginosa*, 168, 169; *incrassata*, 176, 179, 189; *inermis*, 168; *inflata*, 169; *luteofibriata*, 170, 171, 176; *microscopora*, 177; *mutans*, 168; *nigra*, 177; *nigrescens*, 168; *obliqua*, 168; *prunicola*, 168, 169; *punctata*, 169; *sequoiae*, 177, 180; *spiculosa*, 168; *subacida*, 177, 180, 183, 184, 187, 188; *tenuis*, 177, 181, 182, 184, 187, 188; *vaillantii*, 177; *vaporaria*, 177; *weirii*, 168, 177; *xantha*, 177
Portulaca, 22; *cyanosperma*, 24
Posoqueria latifolia, 309
Prosinerpis panamensis, 478
Proteus vulgaris, 158, 161, 163
Prunus avium, 137, 149–156; *mahaleb*, 149; *pennsylvanica*, 149; *persica*, 137–151, 155, 156; *serotina*, 149; *virginiana*, 149
Psychotria brachiata, 309; *bullatifolia*, 309; *cuspidata*, 309; *hartwegiana*, 309; *hirta*, 309; *jamesoniana*, 309; *lateriflora*, 309; *macrophylla*, 309; *patens*, 309; *pilosa*, 309; *recordiana*, 309; *rimbachii*, 309; *rufescens*, 309; *tenuifolia*, 309
Pteridium aquilinum, 402; var. *wightianum*, 422
Pteris aquilina, 422; *biaurita*, 422; var. *quadriaurita*, 422; *calomelanos*, 419; *cretica*, 402, 422; *excelsa*, 422; *farinosa*, 421; *hastata*, 419; *longifolia*, 402, 422; *nitidula*, 419; *quadriaurita*, 422; *stelleri*, 418; *vittata*, 422
Ptilanthus, 472; **scandens**, 472
Ptilidium ciliare, 507, 509; *pulcherrimum*, 509
Ptychogaster cubensis, 168; *rubescens*, 177
Puccinia andropogonicola, 209, 211; *andropogonis-micranthi*, 211; *cacao*, 211; *caracasana*, 213; *cesatii*, 209; *chaetochloae*,

212; *dolosa*, 212; **dolosoides**, 209–212; *incompleta*, 209; **kampalensis**, 211; **kenmorensis**, 209, 211; *minussensis*, 214; var. **africana**, 213; *nakanishikii*, 209; **paspalina**, 211, 212, 217; *penniseti*, 213; *propinqua*, 209; *scleriae*, 213; **stenotaphri**, 213; *substriata*, 212
Pucciniosira deightonii, 218
Pylaisia intricata, 511; *selwynii*, 511
Pythiomorpha gonapodyides, 166
Pythium butleri, 166; *helicoides*, 166

Quercus incana, 406

Radula complanata, 510; *obconica*, 510
Radulum orbiculare, 177, 180, 186, 187
Ravenellii bottomleyae, 215; **hansfordii**, 214, 215; *laevioides*, 215; **maranguensis**, 211, 215
Reboulia hemisphaerica, 511
Relbunium ciliatum, 310; *hirsutum*, 310; *hypocarpeum*, 311; *nitidum*, 311
Remijia pedunculata, 464
Report of some recent collections of Rubiaceae from Ecuador, 295
Rhacomitrium heterostichum var. *gracilescens*, 508, 513
Rhabdoweisia denticulata, 512
Rhizidiomyces apophysatus, 47, 48; *bivellatus*, 47, 48, 50; *hansonae*, 47, 48; **hirsutus** sp. nov., a hairy anisochytrid from Brazil, 47
Rhizopus nigricans, 166
Rhodobryum roseum, 513
Rhytidiadelphus triquetrus, 519
Rhytidium rugosum, 519
Ricinus communis, 566
Ribes odoratissimum, 263
RICKETT, H. W., New combinations in *Cornus*, 223
RILEY, HERBERT PARKES, Inheritance of orange and yellow flower color in *Nemesia strumosa*, 435
ROBBINS, WILLIAM, C., A survey of some wood-destroying and other fungi for antibacterial activity, 165
ROBBINS, WILLIAM J., A survey of some wood-destroying and other fungi for antibacterial activity, 165; Effect of cotton on the germination of phycomyces spores, 76
Rondeletia trichotoma, 311
Roraimanthus imthurnianus, 528
Rubiaceae, report of some recent collections from Ecuador, 295
Rubus odoratus, 349
Rudgea cornifolia, 311; *fimbriata*, 311
Rusts, tropical, 205–222

Sabicea paraensis, 311; *villosa*, 311
Salvinia natans, 424
Sapota sideroxylon, 556
Sapotaceae, studies in, 191, 550
Sarcina lutea, 158–161, 163
Sauvagesia, 521–540; *adima*, 528, 529; *amoena*, 526, 534, 535, 538; *angustifolia*, 522, 532; *brownei*, 523, 525, 527; *deficiens*, 522, 526, 536, 537; *deflexifolia*, 526, 538, 539; *duckei*, 533; *elata*, 526, 531; *erecta*, 523, 525–532; *fruticosa*, 522, 532; *geminiflora*, 529; *gracilis*, 539; **grandifolia**, 526, 530; *imthurniana*, 526, 528, 539; *inconspicua*, 539; *kapleri*, 532; *laxa*, 529; *linearifolia*, 526, 537, 538; *longifolia*, 524, 526, 534; *microphylla*, 526, 527; *nana*, 539; *nutans*, 528; *peruviana*, 528; *pulchella*, 526, 539; *pusilla*, 537; *racemosa*, 526, 532; *ramosissima*, 526, 535; *rosacea*, 532, 533; *rubiginosa*, 529; *serpyllifolia*, 532; **smithiana**, 522, 524, 526, 536, 537; *sprengelii*, 522, 523, 526, 532, 534; *stenophylla*, 526, 527; *striata*, 533; *tenella*, 523, 524, 526, 539
Schizophyllum commune, 166, 168
SCHMITT, MARY BARTLEY, Effect of cotton on the germination of *Phycomyces* spores, 76
SCHNEIDER, HENRY, The anatomy of peach and cherry phloem, 137
SCHNOOBERGER, IRMA, The bryophytes of Shenandoah National Park, Virginia, 506
Schwetschkeopsis denticulata, 515
Sciaromium lescurii, 516
Scilla campanulata, 573
Scleria melanomphala, 218; *naumanniana*, 213
Sclerotium delphinii, 166; *rolfsii*, 166
Scopania undulata, 510
Sedum beyrichianum, 376; *nevei*, 376; *nuttallianum*, 375; *pulchellum*, 375–377; *ternatum*, 376
Selaginella aitchisoni, 426; *borealis*, 426; *jacquemontii*, 426; *sanguinolenta*, 426
Sematophyllum caespitosum, 508, 518; *carolinianum*, 518; *marylandicum*, 518
Serratia marcescens, 158, 161
Setaria aequalis, 206; *pallida-fusca*, 207; *rubiginosa*, 207; *verticillata*, 207
Shaferodendron mayarensis, 558; *moaensis*, 558, 559
Shoot of *Botrychium* interpreted as a series of dichotomies, 491

Sloanea emarginata, 557
Solanum anomalum, 212
Solidago, genes in, 235–244; *rugosa*, 237, 238; var. *typica*, 237, 243; *sempervirens*, 237–243; subsp. *mexicana*, 241; subsp. *typica*, 241; var. *mexicana*, 237, 239, 243; var. *typica*, 237, 239, 243
Some Melastomaceae of Colombia, 472
Sparassis crispa, 177
Sphaerophorus globosus, 163
Sphaerophragmium acaciae, 216
Sphenospora mera, 214
Sporotrichum globulum, 166
Staphylococcus albus, 158, 161; *aureus*, 158, 160–189
ST. JOHN, HAROLD, *Dryopteris, Deschampsia, Portulaca, Lupinus, Fagara, Stenogyne*, and *Dubautia*. Hawaiian plant studies 12, 22
STEERE, WILLIAM CAMPBELL, A report of some recent collections of Rubiaceae from Ecuador, 295; The discovery and distribution of *Cinchona pitayensis* in Ecuador, 464
Stenogyne, 22; *kaalae*, 25; var. *coriacea*, 25; *mollis*, 25; *rugosa*, 25, 26; var. *mollis*, 25; var. *subulata*, 25
Stenotaphrum dimidiatum, 213
Stereocaulon paschale, 161
Stereum albobadium, 168; *fasciatum*, 177; *frustulosum*, 169; *fuscum*, 177; *gausapatum*, 177, 180, 186; *hirsutum*, 168; *ochraceoflavum*, 168; *murrayi*, 166, 179, 180; *pini*, 168; *purpureum*, 168; *rameale*, 169; *roseo-carnum*, 168; *rufum*, 169; *rugosum*, 178, 180, 187; *sanguinolentum*, 168, 169; *sericeum*, 169; *spadiceum*, 168; *subpileatum*, 178; *sulcatum*, 178, 180, 187
STEWART, RALPH R., The ferns of Kashmir, 399
Streptococcus haemolyticus, 158, 161; *viridans*, 158, 161
Studies in the Sapotaceae—I. The North American species of *Chrysophyllum*, 191; IV. The North American species of *Manilkara*, 550
Survey of some wood-destroying and other fungi for antibacterial activity, 165
Symphoricarpos leucocarpus, 263
Syngramme fraxinea, 419; *vestita*, 418

TAFT, CLARENCE E., The validity of the algal genus *Thamniastrum*, 246
Taxonomy of the genus *Sauvagesia* (Ochnaceae), 521
Tetraphis pellucida, 513
Thamniastrum, the genus, 246
Thamnolia vermicularis, 163
Thelia asprella, 515; *hirtella*, 515
Thomas Jefferson, his interest in plant life as revealed in his writings—II, 248
Thuidium abietinum, 515; *delicatulum*, 515; *philiberti*, 508, 515; *recognitum*, 515; *scitum*, 515; *virginianum*, 516
Tobacco, glandular leaf-hairs of, 345
Todea barbara, 500
Topobea, 385; *alternifolia*, 390; **barbata**, 393; **longisepala**, 392; **pubescens**, 392; *stephanochaeta*, 392
Tortella caespitosa, 512
Trametes alaskanae, 178; *americana*, 178; *carnea*, 178; *cubensis*, 168; *heteromorpha*, 178; *hispida*, 168; *malicola*, 178; *rigida*, 168; *rubescens*, 178, 180, 187; *sepium*, 178; *serialis*, 178; *suaveolens*, 168
Trichophyton crateriforme, 171; *gypseum*, 166; *interdigitale*, 171; *mentagrophytes*, 171; *purpureum*, 178; *rosaceum*, 171; *tonsurans*, 171; *violaceum*, 171
Tropical rusts, 205–222
Tyleria floribunda, 523

Ulota americana, 514; *crispa*, 514
Umbilicaria dillenii, 163; *papulosa*, 163
Uredo achyranthicola, 219; achyranthis, 219; **arundinellae-nepalensis**, 218; *brachylaenae*, 221; **corbiculoides**, 206, 211, 219, 220; **crassiperidiata**, 218; **diocleae**, 219; *elephantopodis*, 221; *famelica*, 219; *gossypii*, 206; *grewii*, 220; *longaensis*, 220; *maranguensis*, 215, 216; *melinidis*, 207; *mira*, 211; **palpigera**, 221; *paspalina*, 212; **pergulariae**, 220; **perscita**, 220, 221; *stenoptaphri*, 213; *stereospermi*, 205; *toroiana*, 220, 221; *vernonicola*, 220, 221
Uromyces aspiliicola, 208, 217; *baccarinii*, 208; *leptodermus*, 207; *pretoriensis*, 208; *schinzianus*, 208, 211
Urostigma altissimum, 394, 396; *lacciferum*, 294, 397
Urtica dioica, 349
Usnea florida, 161, 164

Vaccinium hybrids and the development of new horticultural material, 1; *altomontanum*, 3, 7, 16; *amoenum*, 3, 8; *angustifolium*, 3, 6, 11, 18; *arctostaphylos*, 3; *arkansanum*, 3, 7, 8; *ashei*, 3, 4, 8, 16–19; *atrococcum*, 2, 4–6, 11, 19, 46; *australe*, 3, 4, 6, 7, 8, 19; *brittonii*, 3, 6, 7, 11; *caesariense*, 5, 6; *constablaei*, 3, 4; *corymbosum*, 3, 6, 7, 9, 11; *cuthbertii*, 5; *darrowi*, 2, 4, 5, 12; *elliottii*, 3–6, 12;

fuscatum, 8; *hirsutum*, 3; *lamarckii*, 3, 4, 6, 7, 11; *macrocarpum*, 1; *margarettae*, 5; *marianum*, 6, 7; *missouriense*, 5; *myrsinites*, 3, 4, 7, 8, 18, 19; *myrtilloides*, 2, 3, 6, 16, 17; *ovatum*, 3; *pallidum*, 2, 3, 5, 18; *simulatum*, 3, 7; *tallapusae*, 3, 4, 7, 18; *tenellum*, 2–5, 8, 12; *torreyanum*, 46; *uliginosum*, 1; *vacillans*, 5, 6, 11; var. *crinitum*, 5; *virgatum*, 3, 4, 8, 16

Validity of the algal genus *Thamniastrum*, 246

Variations and origin of *Botrychium lanuginosum*, 454

Varinga latifolia, 394, 396, 398

Vernonia amygdalina, 220

Verticillium buxi, 168

Vicia faba, 569, 570

Volutella buxi, 168

Volvox, morphological and cytological study of a new form of,—I, 86; II, 121; *africanus*, 89, 90, 132; *aureus*, 88, 89, 104, 106, 121, 125, 128; *capensis*, 89, 107; *carteri*, 86–113, 121–134; var. **hazeni**, 111, 129, 130, 132, 134; var. **homothallicus**, 132, 134; var. *weismannia*, 128; f. *magariensis*, 129; *gigas*, 89, 99, 132; *globator*, 88, 89, 104, 128; *migulae*, 132, 134; *minor*, 88, 121; *mononae*, 128, 132, 134; *obversus*, 131, 132; *perglobator*, 104, 128; *rousseletii*, 89, 106, 107, 113; *spermatosphaera*, 128; *tertius*, 89–91, 99, 104, 106, 129, 132, 133; *weismannia*, 128, 129, 132

Wedelia oblonga, 213

Weisia viridula, 512

WOLF, F. A., Glandular leaf-hairs of Oriental tobacco, 345

Wood-destroying fungi, 166–189

Woodsia alpina, 402, 404; *hyperborea*, 404

Woodwardia radicans, 401, 408, 418

WYNNE, FRANCES E., The bryophytes of Shenandoah National Park, Virginia, 506

X-rays, effect of on plant tissue, 566, 567, 572, 573

Xanthoria parietina, 163, 164

Xanthoxylum dipetalum var. *geminicarpum*, 24; *mauiense* var. *anceps* f. *petiolulatum*, 25

Zimapania schiedeana, 37